

# *Mycorrhizae for a sustainable world*

Article

Accepted Version

Pickles, B. J., Truong, C., Watts-Williams, S. J. and Bueno, C. G. (2020) Mycorrhizae for a sustainable world. *New Phytologist*, 225 (3). pp. 1065-1069. ISSN 1469-8137 doi: <https://doi.org/10.1111/nph.16307> Available at <https://centaur.reading.ac.uk/87160/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/nph.16307>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# **Mycorrhizae for a sustainable world**

**The 10<sup>th</sup> International Conference on Mycorrhiza (ICOM10), Mérida, Mexico, June 30 – July 5 2019**

Author list: Brian J. Pickles<sup>1</sup>, Camille Truong<sup>2</sup>, Stephanie Watts-Williams<sup>3</sup>, C. Guillermo Bueno<sup>4</sup>

Author affiliations: <sup>1</sup>School of Biological Sciences, University of Reading, Harborne Building, Whiteknights, Reading, RG6 8AS, UK. <sup>2</sup>Instituto de Biología, Universidad Nacional Autónoma de México, Circuito Zona Deportiva s.n., Ciudad Universitaria, Apartado Postal 70-367, C.P. 04510, Ciudad de México, Mexico. <sup>3</sup>School of Agriculture, Food and Wine, University of Adelaide, Glen Osmond, SA, 5064, Australia. <sup>4</sup>Department of Botany, Institute of Ecology and Earth Sciences, University of Tartu, 40 Lai Street, 51005 Tartu, Estonia.

ORCID: <sup>1</sup><https://orcid.org/0000-0002-9809-6455>. <sup>2</sup><https://orcid.org/0000-0002-8510-1761>. <sup>3</sup><https://orcid.org/0000-0003-3467-0662>. <sup>4</sup><https://orcid.org/0000-0002-7288-2271>.

Author for correspondence:

***Brian J Pickles***

Email: [b.j.pickles@reading.ac.uk](mailto:b.j.pickles@reading.ac.uk)

Tel: +44 (0) 118 378 7955

Word count – Text: 1868, Acknowledgements: 120.

Figures: 2 [Colour figures: 2]

Key words: mycorrhiza, agricultural sustainability, ecosystem restoration, arbuscular mycorrhiza, conservation

Conflict of interest statement: The authors declare no conflict of interest.

## Meeting report

More than 80% of plant species exchange resources with mycorrhizal fungi and these associations impact both partners at multiple scales, from individuals to ecosystems. 172 participants from 33 countries and 160 institutions met at the 10<sup>th</sup> International Conference on Mycorrhiza in the city of Mérida in the Yucatan peninsula in Mexico – an area famous for its Mayan archaeological sites, cenotes, and the Chicxulub impact crater that marks the end of the Cretaceous period. They discussed latest advances on mycorrhizal research across 125 talks and 111 posters in 14 sessions focused on the biology, physiology, ecology, evolution and conservation of these interactions from molecules to biomes (Fig. 1). In particular, the contribution of mycorrhizal research to sustainability in agriculture, conservation and ecosystem restoration (Fig. 2) emerged as a promising topic to address today's challenges in the realm of human population growth, globalization and climate change.

### 1. Sustainability in agriculture (managed ecosystems)

Several speakers discussed the increasing abundance of commercial arbuscular mycorrhizal products for agriculture, from “biofertilisers” to advances in seed coating technology (i.e. adding mycorrhizal fungal spores directly to seeds along with nutrients and plant-helper bacteria), and the interest in these products from growers.

A noticeable recurring theme was that commercial biofertilisers make rather dramatic claims about their effectiveness, without evidence of their application leading to direct improvements in crop yield or nutrition. Jan Jansa (Czech Academy of Sciences, Prague, Czech Rep.)'s keynote presentation made the point that biofertilisers do not *create* nutrients per se; however, they may help plants to access existing nutrient sources, and may provide non-nutritional benefits. Examples of non-nutritional mycorrhizal benefits include increased soil glomalin inputs, tolerance of microplastics inputs, and alteration of the soil microbiome (Svenningsen *et al.*, 2018; de Souza Machado *et al.*, 2019; Hestrin *et al.*, 2019). Marcel van der Heijden (Agroscope, Zurich, Switzerland) and Ashleigh Elliott (University of Leeds, Leeds, UK) presented data on the application of commercial inoculants in field and glasshouse trials; crops grown with a commercial arbuscular mycorrhizal (AM) fungal inoculant exhibited higher root colonisation but there were few benefits to growth. Notably, the quality (in terms of active AM fungal propagules) and effectiveness of different commercial products was highly variable.

Miranda Hart (University of British Columbia, Kelowna, Canada) made the case that the variable responses of AM fungal inoculum in field trials are like those observed in the case of plant species invasions (Thomsen & Hart, 2018), and that current practices were too focused on establishing the most vigorous AM fungi. Some important questions arose from the workshop discussion on the topic: how can we, as a research community, contribute to ensuring that mycorrhizal fungal inoculum products are a) appropriate, e.g. are we selecting the most suitable fungi for a given system, rather than good invaders?; and b) successful, e.g. would a “certificate of effectiveness” be required?

In terms of alternative approaches to agriculture, Rillig & Lehmann (2019) identified approximately 285,000 combinations of agricultural practices. In his keynote, Jan Jansa emphasised the need to rigorously quantify the AM symbiosis and its effects in the field, to enable the production of equations and models that make useful predictions, so that we can best make use of the AM symbiosis as a valuable biological resource. Jansa also highlighted the potential to look further down the production chain not only to crop productivity but to the quality of the food product (e.g. do mycorrhizas affect food nutritional content, taste, or spoilage), such as lowered pest impacts, postharvest disease reduction, and thus reduction of food waste (e.g. AM fungi for food security).

## **2. Sustainability in conservation and restoration (natural ecosystems)**

Effective use of the mycorrhizal symbiosis for restoration and conservation requires a deeper understanding of mycorrhizal functionality and related ecosystem processes, and how these processes and functions are altered through interactions with other actors and changing conditions. For instance, several talks (e.g. Heike Bücking, South Dakota State University, Brookings, USA; Ricardo Arraiano Castilho, Kew Gardens, London, UK) highlighted the importance of local soil factors and host nutrient demand in shaping mycorrhizal fungal communities, and whether changes in local environmental conditions associated with climate change (i.e. drought) or nutrient deposition (i.e. soil fertility) may disrupt the structure of these communities. Many speakers discussed the contribution of mycorrhizas in low impact, sustainable approaches to ecosystem restoration (e.g. Brian Pickles, University of Reading, Reading, UK; Cameron Egan, University of Hawai‘i, Mānoa, USA) and species conservation (e.g. Nicole Hynson, University of Hawai‘i, Mānoa, USA; Louise Egerton-Warburton, Chicago Botanic Garden, Glencoe, USA). Still, other mechanisms related to the activities of

mycorrhizal fungi, such as carbon sequestration (i.e. priming effect discussed by María Pozo, EEZ-CSIC, Granada, Spain; and Johanna Pausch, University of Bayreuth, Bayreuth, Germany) or the outcome of interactions among important actors, such as signaling pathways for kin recognition (e.g. Monika Gorzelak, Agriculture and Agri-Food Canada, Lethbridge, Canada), need to be accounted for when considering mycorrhizal applications in conservation and ecosystem restoration.

A key theme is that different ecosystems may well need different approaches (i.e. there is no “silver bullet” for restoration or conservation). For example, Louise Egerton-Warburton found that “cedar” (*Widdringtonia wrighteii*) seedlings grew well in nursery conditions but experienced drastic mortality following transfer to the field in Malawi. In turn, Nicole Hynson and Cameron Egan’s work showed that incomplete recovery of Hawaiian native fungal communities following successful growth of planted native host trees may compromise forest restoration. However, the presence of diverse mycorrhizal fungal communities is not the only requirement for a successful restoration plan. For instance, when comparing the performance of AM fungal species on high- and low-quality (determined by associated fungal biomass) native plant hosts in tallgrass prairie, Ylva Lekberg (MPG Ranch, Missoula, USA) found that AM fungal identity and abundance influenced plant performance, while AM fungal species diversity was unimportant in this regard. Similar results were found in a successional plant-feedback study where only the appropriate late successional AM fungi with their corresponding plant species grew faster and larger (Kozioł & Bever, 2019). In the North American Southwest, Catherine (Kitty) Gehring (Northern Arizona University, Flagstaff, USA) found that intraspecific drought tolerance of pinyon pine was strongly associated with root-colonising ECM fungal species composition. Here, drought-tolerant pinyons tended to associate with *Geospora* spp., which increased water flow velocity in drought-tolerant seedling lineages and reduced it in intolerant lineages. A related study revealed that after successive droughts, ECM fungal species composition and abundance in roots of pinyon pines were responsive to tree mortality, with *Geospora* increasing and *Tuber* spp. decreasing in response to pine death (Mueller *et al.*, 2019). These studies indicate that the identity of mycorrhizal fungi and their interaction with certain host traits are critical for achieving restoration aims.

### 3. Advances in mycorrhizal research with sustainability applications

Understanding patterns of plant mycorrhizal type dominance, for instance in highly protected and valuable ecosystems, is key to understanding many ecosystem processes and their dynamics, and hence predicting limiting factors and environmental risks. In his keynote presentation, Richard Phillips (Indiana University, Bloomington, USA) presented a plethora of works describing differences in functioning between forests dominated by AM and ECM trees in similar climatic conditions (e.g. Zhang *et al.*, 2018). It had long been hypothesised that ECM-dominated forests accumulate more soil carbon, due in part to visibly greater production of recalcitrant organic matter. Yet when soils from ECM- and AM-dominated forests in proximity were compared to a depth of 1 m, greater accumulation of soil organic matter was found in AM-dominated forests (Craig *et al.*, 2018). Several talks presented at ICOM10 highlighted how processes such as C storage, soil enzymatic activities, nutrient cycling, and ecosystem-level sensitivity to global changes may vary (in part) because of mycorrhizal interactions (e.g. Haley Dunleavy Northern Arizona University, Flagstaff, USA; Tom Thirkell, University of Leeds, Leeds, UK; Melanie Jones, University of British Columbia, Kelowna, Canada). These results clearly stress the need to consider how the dominance of different mycorrhizal types may impact ecosystem function, and the consequences of host changes for broader ecosystem dynamics, management, and restoration. Nonetheless, subdominant plant species such as herbs and grasses in the forest understory can also play significant roles in ecosystems. For example, Rebecca Bunn (Western Washington University, Bellingham, USA) revisited the ‘direct mineral cycling hypothesis’ from the 1960’s and showed that AM fungal hyphae are active in forest leaf litter through cooperation with other microorganisms (e.g. Lin Zhang, China Agricultural University, Beijing, China), even in ecosystems dominated by ECM trees (Bunn *et al.* 2019). Despite these recent advances in using plant mycorrhizal type to investigate ecosystem processes, distinguishing between the plant mycorrhizal types (such as AM, ECM, or dual AM and ECM) is not always easily solved and different approaches coexist (Brundrett & Tedersoo, 2019; Bueno *et al.* 2019). ICOM10 facilitated an interesting debate in this respect, discussing possibilities for merging functional, morphological, and experimental approaches to tackle this important issue.

Studies of the functions of symbioses in the presence of their closest neighbours are also warranted. Marco Cosme (Université Catholique de Louvain, Louvain-la-Neuve, Belgium) illustrated the role that mycorrhizal fungi can play in ‘non-mycorrhizal’ plant functional responses, in which a presumed non-host species (*Arabidopsis thaliana*) in the presence of a mycorrhizal plant (*Medicago truncatula* colonised by the AM fungus *Rhizophagus* sp.)

exhibited root cortex colonisation. No nutrient exchange (via arbuscules) was observed, but the non-host plant exhibited activation of AM fungal-induced resistance to pathogens (Fernández *et al.*, 2019), indicating a functionally beneficial colonisation of the presumed non-host species. All in all, examining the multifunctional effects of the entire root mycobiome, including non-mycorrhizal and “fine root endophyte” fungi (Hoysted *et al.*, 2019) across plants, may be crucial to predicting the effect of global changes in natural and managed ecosystems.

#### 4. Challenges

During the conference, key challenges facing mycorrhizal research (and researchers) in the coming decades were addressed:

##### *Global change*

Mycorrhizal symbioses are already highly complex, so how do we decipher mycorrhizal effects in systems subjected to multiple simultaneous pressures? Many speakers discussed mycorrhizal responses to climate change impacts such as drought, fire, and insect outbreaks (e.g. Philip Brailey, University of York, York, UK; Jean Carlos Rodríguez-Ramos, University of Alberta, Edmonton, Canada; Yong Zheng, Fujian Normal University, Fuzhou, China). Restoration of ecosystems exposed to pollutants (e.g. microplastics) was another common theme, as exemplified by Matthias Rillig (Freie Universität Berlin, Berlin, Germany)’s keynote talk. Species introductions of exotic fungi and/or exotic hosts are another important topic that potentially leads to fungal invasions. For example, global patterns in native vs introduced island floras revealed a strong tendency towards introduced mycorrhizal plants compared to non-mycorrhizal natives (Delavaux *et al.*, 2019), with some notable exceptions to the general pattern (e.g. Hawaii). Anne Pringle (University of Wisconsin-Madison, Madison, USA)’s research on *Amanita muscaria* (fly agaric) invasions in North America revealed that the population structure of this invasive fungus differed dramatically compared to its native range. The interactive effects of global change processes on mycorrhizal fungi and their hosts will undoubtedly provoke significant research effort from the mycorrhizal research community.

##### *Methodological issues and advances*

Although this topic is not new, finding ecologically relevant control for, and measurement of, the mycorrhizal status of plants is still controversial. Is “non-mycorrhizal” really an appropriate control condition for plants, given the prevalence of mycorrhizal fungi in natural and

anthropogenic ecosystems (i.e., plants without mycorrhizal symbionts are rare), or would severing/restricting common mycorrhizal networks be more relevant experimental control (e.g. David Johnson, University of Manchester, Manchester, UK)?

The advent of modern high-throughput plant phenotyping systems has allowed us to begin characterising mycorrhizal host plant (shoot) growth responses (positive through to negative) over time (Watts-Williams *et al.*, 2019), rather than just at the harvest time point. This technology will be especially useful when it extends to root phenotyping platforms that allow for high resolution screening, and analysis of the effects of mycorrhizal fungi on root growth and morphology over time.

Several issues remain unresolved among the continual technological advances used for molecular work and interpretation of those data, as sequencing of mycorrhizal fungal communities becomes more commonplace. As Annegret Kohler (INRA, Nancy, France)'s keynote talk asked: What does gene copy number mean in terms of function? What does sequence abundance really mean in terms of species abundances? Many researchers uncritically present sequence abundances from NGS platforms as if they were equivalent to species relative abundances, although the ecological relevance of sequence abundance data needs to be cautiously addressed within the mycorrhizal (Nguyen *et al.*, 2015) and wider microbiome (Gloor *et al.*, 2017) research communities. Clearly, there needs to be more care with the use of metagenomic data and this may prove to be a suitable topic for a discussion session at a future ICOM.

## Acknowledgements

This report is dedicated to the memory of the late Professor Sally E. Smith (1941 – 2019; <https://nph.onlinelibrary.wiley.com/doi/full/10.1111/nph.15569>). Sally's illustrious research career spanned more than 50 years and contributed important findings, particularly on the arbuscular mycorrhizal symbiosis. She inspired many young scientists to pursue careers in plant and mycorrhiza research, and she will be greatly missed by the mycorrhiza research community. Sally received the Eminent Mycorrhiza Researcher Award at ICOM10.

The authors thank the local organising committee, the international scientific committee, the International Mycorrhiza Society, and all the delegates at ICOM10 for many stimulating talks, posters, and discussions throughout the conference. CGB was funded by the Estonian Research



Council (grant IUT 20-28), and the European Regional Development Fund (Centre of Excellence EcolChange).

## References

**Brundrett M, Tedersoo L. 2019.** Misdiagnosis of mycorrhizas and inappropriate recycling of data can lead to false conclusions. *New Phytologist* **221**: 18–24.

**Bueno CG, Aldrich-Wolfe L, Chaudhary VB, Gerz M, Helgason T, Hoeksema JD, Klironomos J, Lekberg Y, Leon D, Maherali H, et al. 2019.** Misdiagnosis and uncritical use of plant mycorrhizal data are not the only elephants in the room. *New Phytologist* doi:10.1111/nph.15976.

**Bunn RA, Simpson DT, Bullington LS, Lekberg Y, Janos DP. 2019.** Revisiting the ‘direct mineral cycling’ hypothesis: arbuscular mycorrhizal fungi colonize leaf litter, but why? *The ISME Journal* **13**: 1891–1898.

**Craig ME, Turner BL, Liang C, Clay K, Johnson DJ, Phillips RP. 2018.** Tree mycorrhizal type predicts within-site variability in the storage and distribution of soil organic matter. *Global Change Biology* **24**: 3317–3330.

**Delavaux CS, Weigelt P, Dawson W, Duchicela J, Essl F, van Kleunen M, König C, Pergl J, Pyšek P, Stein A, et al. 2019.** Mycorrhizal fungi influence global plant biogeography. *Nature Ecology & Evolution* **3**: 424–429.

**Fernández I, Cosme M, Stringlis IA, Yu K, Jonge R, van Wees SM, Pozo MJ, Pieterse CMJ, van der Heijden MGA. 2019.** Molecular dialogue between arbuscular mycorrhizal fungi and the nonhost plant *Arabidopsis thaliana* switches from initial detection to antagonism. *New Phytologist* **223**: 867–881.

**Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. 2017.** Microbiome Datasets Are Compositional: And This Is Not Optional. *Frontiers in Microbiology* **8**: article 2224.

**Hestrin R, Hammer EC, Mueller CW, Lehmann J. 2019.** Synergies between mycorrhizal fungi and soil microbial communities increase plant nitrogen acquisition. *Communications Biology* **2**: 233.

**Hoysted G, Jacob A, Kowal J, Giesemann P, Bidartondo M, Duckett J, Gebauer G, Rimington W, Schornack S, Pressel S, et al. 2019.** Mucoromycotina fine root endophyte fungi form nutritional mutualisms with vascular plants. *Plant Physiology*: pp.00729.2019.

**Koziol L, Bever JD. 2019.** Mycorrhizal feedbacks generate positive frequency dependence accelerating grassland succession (B Turner, Ed.). *Journal of Ecology* **107**: 622–632.

**Mueller RC, Scudder CM, Whitham TG, Gehring CA. 2019.** Legacy effects of tree


- mortality mediated by ectomycorrhizal fungal communities. *New Phytologist* **224**: 155–165.
- Nguyen NH, Smith D, Peay K, Kennedy P. 2015.** Parsing ecological signal from noise in next generation amplicon sequencing. *New Phytologist* **205**: 1389–1393.
- Rillig MC, Lehmann A. 2019.** Exploring the agricultural parameter space for crop yield and sustainability. *New Phytologist* **223**: 517–519.
- de Souza Machado AA, Lau CW, Kloas W, Bergmann J, Bachelier JB, Faltin E, Becker R, Görlich AS, Rillig MC. 2019.** Microplastics Can Change Soil Properties and Affect Plant Performance. *Environmental Science & Technology* **53**: 6044–6052.
- Svenningsen NB, Watts-Williams SJ, Joner EJ, Battini F, Efthymiou A, Cruz-Paredes C, Nybroe O, Jakobsen I. 2018.** Suppression of the activity of arbuscular mycorrhizal fungi by the soil microbiota. *The ISME Journal* **12**: 1296–1307.
- Thomsen CN, Hart MM. 2018.** Using invasion theory to predict the fate of arbuscular mycorrhizal fungal inoculants. *Biological Invasions* **20**: 2695–2706.
- Watts-Williams SJ, Jewell N, Brien C, Berger B, Garnett T, Cavagnaro TR. 2019.** Using High-Throughput Phenotyping to Explore Growth Responses to Mycorrhizal Fungi and Zinc in Three Plant Species. *Plant Phenomics* **2019**: 1–12.
- Zhang H-Y, Lü X-T, Hartmann H, Keller A, Han X-G, Trumbore S, Phillips RP. 2018.** Foliar nutrient resorption differs between arbuscular mycorrhizal and ectomycorrhizal trees at local and global scales. *Global Ecology and Biogeography* **27**: 875–885.

287 **Figure legends**


288 **Figure 1.** ICOM10 covered a variety of recent basic and applied mycorrhizal research with a  
289 focus on topics that inform the sustainability of managed and natural ecosystems. Interactions  
290 between global change processes, and the interpretation of data from rapidly advancing  
291 sequencing technologies, emerged as common challenges for mycorrhizal researchers.

292 **Figure 2.** Planned and unplanned (in some cases unwanted) inputs into managed (e.g.,  
293 agricultural, silvicultural) and natural mycorrhizal systems, and potential or existing outputs,  
294 which can extend to ecosystem and socio-economic impacts.

295



**Figure 1.**



**Figure 2.**